

ZADITEN[®] Eye Drops 0.025%

ketotifen

Multidose bottle

You should ensure that you speak to your pharmacist or doctor to obtain the most up to date information on the medicine. You can also download the most up to date Consumer Medicine Information (CMI) leaflet from zaditen.com.au in Australia.

23391900



How to use ZADITEN[®]

Follow all directions given to you by your pharmacist or doctor carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the label, ask your pharmacist or doctor for help.

If you wear soft contact lenses, remove them before using this medicine. Wait for 15 minutes after using it before reinserting your contact lenses.

The ingredients in Zaditen can affect contact lenses.

Follow these steps to use Zaditen multi-dose bottle:

1. Wash your hands thoroughly with soap and water.
2. Open the multi-dose bottle. If the tamper evident snap collar is loose on the multidose bottle, remove before using Zaditen.
3. Make sure the tip of the bottle does not touch anything as this may contaminate the solution.
4. Hold the bottle in one hand between the thumb and forefinger.
5. With your head tilted back, use the forefinger of the other hand to pull down the lower lid of the eye to be treated.
6. Place the dropper close to your eye but without touching your eye or eyelid, and squeeze the bottle gently so that one drop falls into the pouch between your lower lid and your eye, and then release your lower lid.
Only one drop is needed.
7. Close your eye and gently press the inner corner of your eye with your forefinger for one minute.
8. Treat your other eye in the same way if it is also affected.
9. Close the bottle and keep it at room temperature (below 25°C).

Discard the multi-dose bottle and remaining solution four (4) weeks after opening.

If you are using any other eye drops at the same time as Zaditen, leave an interval of at least 5 minutes between each drop to stop the drops being washed out of the eye.

How much to use

The usual dose of Zaditen Eye Drops is one drop in the affected eye(s) twice a day.

ZADCM/1
R08

Care
Pharmaceuticals



23386600



Things to be careful of

Be careful driving or operating machinery until you know how Zaditen affects you.

This medicine may cause blurred vision or tiredness in some people. If you have either of these symptoms, do not drive, operate machinery or do anything else that could be dangerous. Children should be careful when riding bicycles or climbing trees.

Supplied by

In Australia this product is supplied by:

Care Pharmaceuticals Pty Ltd,
Bondi Junction NSW 2022,
Australia
FREECALL Australia 1800 788 870

This leaflet was prepared in April 2022

® = Registered Trademark

AUST R 99979



23386600

ZADCMI/1
R08
Care
Pharmaceuticals



APPLICATION TO REAR PAGE OF PIL/CMII